

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Reg4 in direct ELISAs and Western blots. In direct ELISAs, less than 20% cross-reactivity with recombinant mouse (rm) Reg4, rmReg3A, and recombinant rat (rr) Reg3A is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Reg4 Asp23-Pro158 Accession # Q9BYZ8
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Reg4 (regenerating islet-derived protein 4; also named Reg-like protein, RELP, and gastrointestinal secretory protein) is a 17 kDa, secreted glycoprotein belonging to the calcium (C-type) dependent lectin superfamily and the regenerating gene family (1-3). There are three potential isoforms for human Reg4. Isoform 1 is synthesized as a 158 amino acid (aa) precursor with a 22 aa signal sequence and a 136 aa mature chain. Amino acids 30-156 constitute a C-type lectin-like domain, and amino acid 50 is a site for potential N-linked glycosylation. In addition, amino acids 127 and 142-143 form a ligand-binding surface. In isoform 2, there is a 79 aa substitution for the C-terminal 101 amino acids, while in isoform 3, there is an 11 aa substitution for the C-terminal 57 amino acids. Human Reg4 shares 68% and 67% aa sequence identity with rat and mouse Reg4, respectively. Like other members of the regenerating gene family, Reg4 is expressed in the gastrointestinal (GI) tract and ectopically at other sites in the setting of tissue injury (1, 2). In particular, Reg4 is expressed in normal colon mucosa, and up-regulated in colon adenocarcinoma, pancreatic cancer, gastric adenocarcinoma, inflammatory bowel disease (Crohn's disease and ulcerative colitis), and, outside the GI tract, in prostate adenocarcinoma (2-8). The physiological function of Reg4 is presently unknown, but it may be involved in inflammatory and metaplastic responses of the GI epithelium (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.