

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human RELT/TNFRSF19L in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) 4-1BB, rhBAFF R, rhCD27, rhTAJ, rhCD30, rhDR3, rhDR6, rhTNF RI, rhTNF RII, rhEDAR, rhFas, rhGITR, rhHVEM, rhNGF R, rhOPG, and recombinant mouse OX40 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human RELT/TNFRSF19L Ser26-Ala160 (Arg127Gly, Arg129Gly) Accession # Q969Z4
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

RELT (Receptor Expressed in Lymphoid Tissues) is a type I transmembrane glycoprotein belonging to the tumor necrosis factor receptor superfamily (TNFRSF) and has been designated TNFRSF19-like (TNFRSF19L) (1, 2). It is primarily expressed in hematopoietic tissues and peripheral blood leukocytes. Human RELT cDNA encodes a 430 amino acid (aa) residue precursor protein with a putative 26 aa signal peptide, a 136 aa extracellular domain containing one TNF receptor cysteine-rich domain and one potential N-linked glycosylation site, a 21 aa transmembrane domain and a 247 aa cytoplasmic region containing no death domain. Human RELT shares 85% and 96% aa sequence homology with mouse RELT (Accession # BAC40459) and macaque RELT (Accession # Q9N092), respectively. Among TNFRSF members, the RELT extracellular domain is most closely related to that of TNFRSF19 and OX40. RELT has been shown to exclusively bind the adaptor protein TNF receptor-associated factor 1 (TRAF1). However, it has also been shown to activate the NF-κB pathway independently of TRAFs. Immobilized RELT can co-stimulate T-cell proliferation in the presence of CD3 signaling, suggesting a potential regulatory role in immune response.

### References:

1. <http://www.gene.ucl.ac.uk/nomenclature/genefamily/tnftop.html>.
2. Sica, G. *et al.* (2001) *Blood* **97**:2702.