

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat FGF-BP in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human FGF-BP is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat FGF-BP Glu21-Cys238 Accession # Q9QY10
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast growth factor binding protein (FGF-BP), also known as HBp17, is a secreted glycoprotein that increases the bioavailability of FGFs (1). Mature FGF-BP is a 34 kDa O-glycosylated protein with five conserved intrachain disulfide bonds (2-4). FGF-BP contains a heparin-binding domain (aa 113-146) and a distinct FGF-binding region (aa 197-238) (5). Mature rat FGF-BP shares 54% and 81% aa sequence identity with mouse and rat FGF-BP, respectively. FGF-BP is expressed throughout development and in adult squamous epithelium (6, 7). It is upregulated in injured skin, renal tubular epithelium, and spinal nerves as well as in carcinomas of the skin, colon, and pancreas (3, 8-11). FGF-BP binds FGF -1, -2, -7, -10, and -22 which are secreted and sequestered in the extracellular matrix (ECM) (8, 12). The interactions of FGF-BP with heparin sulfate proteoglycans (HSPG) and FGF, modulates their activities (7, 9, 13, 14). FGF-BP enhances the mitogenic effects of FGFs, thereby contributing to epithelial, endothelial, and neuronal tissue repair, angiogenesis, and tumor growth (8-10, 12, 15, 16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.