

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human DcR3/TNFRSF6B in ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) TRAIL R1, rhTRAIL R2, rhTRAIL R3, and rhTRAIL R4 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human DcR3/TNFRSF6B Val24-His300 Accession # O95407
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human decoy receptor 3 (DcR3), also called TNFRSF6B, TR6, and M68, is a member of the TNF receptor superfamily. The cDNA of DcR3 encodes a 300 amino acid (aa) polypeptide with a putative 23 aa signal peptide. Like osteoprotegerin (OPG), DcR3 lacks a transmembrane sequence and is a secreted protein. DcR3 shares sequence identity with OPG (31%), TNF RII (29%), and Fas (17%). It was found to be expressed in a variety of different tissues and at high levels in many malignant tumors. Among TNF family members, DcR3 was shown to bind with Fas ligand (FasL) and LIGHT and inhibit FasL- and LIGHT-induced apoptosis. Thus, in addition to DcR1, DcR2, and OPG, DcR3 is another TNFR family molecule which modulates ligands that induce apoptosis. Over-expression of DcR3 might be a mechanism by which certain tumors escape immune-cytotoxic attack.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.