

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LRP-6 in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant mouse LRP-6 is observed, and less than 1% cross-reactivity with recombinant human (rh) LRP-1, rhLRP-4, and rhLRP-5 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LRP-6 Ala20-Pro1368 Accession # O75581
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The low-density lipoprotein (LDL) receptor-related protein 5 (LRP-5) and LRP-6 constitute a distinct subgroup of the LDL receptor family. Both LRP-5 and LRP-6 are type I transmembrane proteins that function as co-receptors with Frizzled (FZD) in the canonical Wnt signaling pathway (1, 2). LRP-6 cDNA encodes a 1613 amino acid residue (aa) precursor with a 19 aa signal sequence, a 1353 aa extracellular region, a 23 aa transmembrane (TM) segment and a 218 aa cytoplasmic domain (3). The extracellular region contains four N-terminal cysteine-rich EGF-like repeats, followed by three cysteine-rich LDLR repeats. This pattern of the EGF and LDLR repeat arrangement is different than that typically found in other LDL receptor family proteins. The intracellular region of LRP6 contains protein-protein interaction motifs and is required for canonical Wnt signal transduction (4). Human LRP-6 shares 98% and 74% aa sequence identity with mouse LRP-6 and human LRP-5, respectively. Based on the current model, canonical Wnt signaling requires the interaction of Wnt with FZD and LRP to form a trimeric complex which signals downstream to stabilize cytoplasmic β -catenin. The stabilized β -catenin is then translocated to the nucleus where it complexes with the transcription factor LEF/TCF to regulate the transcription of target genes (5). LRP-6 has also been shown to interact with the Dickkopf proteins (DKKs), which are modulators of Wnt signaling (6-8). Binding of DKK-1 to LRP-6 dissociates LRP-6 from FZD, and antagonizes the formation of the functional receptor complex. On cells where the transmembrane proteins Kremens are also present, a ternary complex of LRP-6, DKK-1 and Kremen is formed to trigger the internalization of the complex and removal LRP6 from the cell surface. Thus, DKK-1 and Kremen function synergistically to antagonize LRP-6-mediated Wnt activity. Although DKK-2 also functions as a Wnt antagonist on cells that express Kremen, DKK-2 binding to LRP-6 in the absence of Kremen activates rather than inhibits LRP mediated β -catenin signaling (9, 10).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.