

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse Soggy-1/DkkL1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human Soggy-1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Soggy-1/DkkL1 Leu21-Leu230 Accession # AAF02679
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Soggy-1, also known as DkkL1 (Dickkopf-like 1), is a secreted protein that is related to the Dickkopf family of Wnt antagonists. Dkk-1, -2, -3, and -4 each have two cysteine-rich domains separated by a linker. Soggy-1 does not contain cysteine-rich domains but bears some sequence homology with the N-terminal regions of Dkk proteins (1, 2). The mouse Soggy-1 cDNA encodes a 230 amino acid (aa) precursor that includes a 20 aa signal sequence (2). Mouse Soggy-1 shares 65% and 90% aa sequence identity with human and rat Soggy-1, respectively, and approximately 15% aa sequence identity with mouse Dkk-1, -2, -3, and -4. Mouse Soggy-1 is expressed at various sites in the embryo but in the adult is primarily found in the testes (3-5). Soggy-1 transcription is regulated by the spermatocyte specific factor, RFX2 (6). The regulatory elements for Soggy-1 lie very close to those of TEAD-2, a transcription factor that is expressed very early in development (4). Soggy-1 and TEAD-2 are co-expressed in preimplantation embryos and embryonic stem cells, but differentiated cells express only one or the other (4, 5). During development, Soggy-1 is first detectable at the onset of sexual differentiation (4). Soggy-1 is localized to the acrosome in developing spermatocytes and mature spermatozoa (3, 4). Soggy-1, as expressed in developing mouse spermatocytes, is a 34 kDa N-glycosylated protein. This glycosylation is not present on Soggy-1 in mature spermatozoa, although the apparent molecular weight suggests the presence of some post-translational modification (3). Two shorter forms of Soggy-1 have been described that result from the use of internal methionine residues for initiation (4).

References:

1. Kawano, Y. and R. Kypta (2003) *J. Cell Sci.* **116**:2627.
2. Krupnik, V.E. *et al.* (1999) *Gene* **238**:301.
3. Kohn, M.J. *et al.* (2005) *Mol. Reprod. Dev.* **71**:516.
4. Kaneko, K.J. and M.L. DePamphilis (2000) *Nucleic Acids Res.* **28**:3982.
5. Kaneko, K.J. *et al.* (2004) *Mol. Cell. Biol.* **24**:1968.
6. Horvath, G.C. *et al.* (2004) *Biol. Reprod.* **71**:1551.