

## DESCRIPTION





<b>Species Reactivity</b>	Human/Mouse/Rat
<b>Specificity</b>	Detects rat FABP1/L-FABP in direct ELISAs and Western blots. In Western blots, less than 5% cross-reactivity with recombinant rat FABP2, recombinant human FABP3, recombinant mouse (rm) FABP4, and rmFABP5 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant rat FABP1/L-FABP Met1-Ile127 Accession # P02692
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.2-0.5 µg/mL	See Below
<b>Immunocytochemistry</b>	5-15 µg/mL	Immersion fixed rat liver cells
<b>Immunohistochemistry</b>	5-15 µg/mL	See Below
<b>Simple Western</b>	10 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

FABP1, also known as liver FABP (L-FABP, Z-protein and squalene, and sterol-carrier protein [SCP]) is a member of the intracellular FABP family. It is highly expressed in the liver, intestine, kidney, and lung. FABP1 binds free fatty acids, their co-enzyme A derivatives, and may be involved in intracellular lipid transport.