

DESCRIPTION





Species Reactivity	Human/Mouse/Rat
Specificity	Detects rat FABP1/L-FABP in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant rat FABP1/L-FABP Met1-Ile127 Accession # P02692
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.2-0.5 µg/mL	See Below
Immunocytochemistry	5-15 µg/mL	Immersion fixed rat liver cells
Immunohistochemistry	5-15 µg/mL	See Below
Simple Western	10 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

FABP1, also known as liver FABP (L-FABP, Z-protein and squalene, and sterol-carrier protein [SCP]) is a member of the intracellular FABP family. It is highly expressed in the liver, intestine, kidney, and lung. FABP1 binds free fatty acids, their co-enzyme A derivatives, and may be involved in intracellular lipid transport.