

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Fcγ RIIIA/B (CD16) in direct ELISAs and Western blots. In these formats, approximately 5% cross-reactivity with recombinant human Fcγ RIIA is observed and less than 2% cross-reactivity with recombinant mouse CD16 is
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Fcγ RIIIB (R&D Systems, Catalog # 1597-FC) Thr20-Gln208 Accession # O75015
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Receptors for the Fc region of IgG (Fcγ R) are members of the Ig superfamily. Based on their genetic organization and molecular structure, three classes of human Fcγ Rs: RI (CD64), RII (CD32), and RIIIB (CD16), which generate multiple isoforms, are recognized (1 - 3). These receptors function in the activation or inhibition of immune responses. The activating-type receptor either has, or associates non-covalently with an accessory subunit (Fcγ Rγ or ζ chain) that has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. In contrast, the inhibitory receptor (Fcγ RIIIB) has a built-in immunoreceptor tyrosine-based inhibitory motif (ITIM) in its own cytoplasmic domain. Fcγ RI is a high-affinity receptor that binds monomeric IgG. Both Fcγ RII and RIIIB are low-affinity receptors that bind IgG in the form of immune complexes. Two genes for human Fcγ RIIIB, A and B, encoding a transmembrane receptor and a glycosylphosphatidylinositol (GPI) anchored protein, respectively, have been identified. Three allelic variants of Fcγ RIIIB, NA-1, NA-2, and SH, exist. A soluble form of Fcγ RIIIB corresponding to the extracellular region of the receptor is produced by proteolytic cleavage and circulates in plasma and other body fluids. The extracellular domains of Fcγ RIIIA and B share 97% amino acid sequence homology. Whereas Fcγ RIIIA is expressed on most effector cells of the immune system including macrophage, monocyte, NK cells, mast cells, eosinophils, dendritic cells, and Langerhans cells, Fcγ RIIIB is selectively expressed in neutrophils and eosinophils. Signaling through Fcγ RIIIA results in oxidative burst, cytokine release and phagocytosis by macrophages, antibody-dependent cellular cytotoxicity by natural killer cells and degranulation of mast cells. By contrast, Fcγ RIIIB is a decoy receptor that binds IgG complexes without triggering activation. Soluble Fcγ RIIIB has a regulatory role in inflammatory processes (4). It interacts with complement receptors CR3 and CR4 on monocytes to induce the production of pro-inflammatory cytokines.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.