

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse TWEAK R in ELISAs and Western blots. In sandwich ELISAs, less than 2% cross-reactivity with recombinant human TWEAK R is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse TWEAK R/TNFRSF12 Glu28-Trp79 Accession # Q9CR75
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TNF-related weak inducer of apoptosis receptor (TWEAK R) belongs to the TNF receptor superfamily and is designated TNFRSF12. The gene for TWEAK R was originally identified as a fibroblast growth factor-inducible immediate-early response gene Fn14 in mouse NIH 3T3 fibroblasts (1, 2). Mouse TWEAK R cDNA encodes a 129 amino acid (aa) residues type I transmembrane protein with a 27 aa signal peptide, a 53 aa extracellular domain, a 21 aa transmembrane domain, and a 28 aa cytoplasmic domain (1-3). Human and mouse TWEAK R share 82% aa sequence identity. TWEAK R is the smallest member of the TNF receptor superfamily and contains only one cysteine-rich region in its extracellular domain. The TWEAK R cytoplasmic domain contains one TRAF binding motif which binds TRAFs 1, 2, and 3. TWEAK R binds its ligand TWEAK/TNFSF12 with high affinity to initiate a signal transduction cascade that depending upon the cell type, may lead to a variety of cellular responses including cell death by both caspase-dependent apoptosis and cathepsin B-dependent necrosis, cell proliferation, and angiogenesis (2-6). In newborn mice, TWEAK R is highly expressed in all tissues examined (heart, intestine, kidney, liver, lung, and skin) (1). In adult mice, high TWEAK R expression levels are found in the heart and ovary, while lower expression levels are detected in the lung, kidney, skin. Elevated levels of TWEAK R mRNA were found in human or mouse hepatocellular carcinoma specimens, in regenerating mouse liver and in injured rat arteries (2, 3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.