

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Tryptase γ -1/TPSG1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Tryptase γ -1/TPSG1 Arg20-Arg281 Accession # Q9NRR2
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

As mediators of inflammatory and allergic response, mast cells are found throughout the body concentrated near blood vessels in connective tissue and the mucous membranes of the respiratory and gastrointestinal tract (3). Upon activation, human mast cells release granules that are enriched with neutral serine proteases including Tryptases, chymase and cathepsin G (4). Tryptase γ -1, also called transmembrane Tryptase, is encoded by TPSG1, one of many serine protease genes clustered in human chromosome 16p13.3 (5). Human Tryptase γ -1 is synthesized as a 321 amino acid preproenzyme with a C-terminal transmembrane anchor (1, 2). The rhTPSG1 was expressed as a soluble protein terminated at residue 281 and corresponded to the proenzyme. The proenzyme can be cleaved by trypsin to form the active enzyme. The serine protease activity of trypsin-activated rhTPSG1 can be inhibited by ecotin (R&D Systems, Catalog # 1328-PI). Greater than 95% protease activity is inhibited by ecotin at approximately 10:1 molar ratio.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.