

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Jagged 2 in direct ELISAs and Western blots. In Western blots, approximately 35% cross-reactivity with recombinant human Jagged 1 is observed and less than 1% cross-reactivity with recombinant rat Jagged 1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Jagged 2 Met27-Asp307 Accession # Q9Y219
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human Jagged 2 is a 131 kDa (predicted) member of the Delta-Serrate-Lag-2 (DSL) family of ligands. This family activates LIN12/Notch proteins and thereby regulates cell fate determination during development (1-5). It is a type 1 transmembrane protein that is synthesized as a 1238 amino acid (aa) precursor. It contains a 23 aa signal sequence, a large 1057 aa extracellular region, a 21 aa transmembrane region, and a short 137 aa cytoplasmic region. The extracellular region contains four potential N-linked glycosylation sites, a DSL domain, 16 EGF-like repeats (many of which are also sites of calcium binding), a von Willebrand factor (vWF) type C domain, and a cysteine-rich region just proximal to the transmembrane segment (2). There are two isoforms for human Jagged 2, named long and short. The short form lacks a splicing variant region (aa 421-461) that is present in the long form of the protein. Human Jagged 2 shares 90% and 87% aa sequence identity with mouse and rat Jagged 2, respectively. During murine embryonic development, Jagged 2 is expressed highest in fetal thymus, epidermis, foregut, dorsal root ganglia, and inner ear (2). In 2 week old mice, the Jagged 2 transcript is most abundant in heart, lung, thymus, skeletal muscle, brain, and testis (2). Functionally, it is suggested that Jagged 2 engages the Notch1 pathway of signal transduction (2). It is involved in the development of the mammalian limb, branchial arches, central and peripheral nervous systems, T cell lineage differentiation, natural killer cells, and the establishment of functional natural killer cell lines (3, 5, 6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.