

#### DESCRIPTION

|                           |                                                                                                                                                                                                                                                                             |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Canine                                                                                                                                                                                                                                                                      |
| <b>Specificity</b>        | Detects canine CCL2/JE/MCP-1 in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant human (rh) CCL2/MCP-1 is observed, approximately 10% cross-reactivity with recombinant mouse (rm) JE, recombin                       |
| <b>Source</b>             | Polyclonal Goat IgG                                                                                                                                                                                                                                                         |
| <b>Purification</b>       | Antigen Affinity-purified                                                                                                                                                                                                                                                   |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant canine CCL2/JE/MCP-1<br>Gln24-Pro101<br>Accession # P52203                                                                                                                                                                              |
| <b>Conjugate</b>          | Alexa Fluor 647<br>Excitation Wavelength: 650 nm<br>Emission Wavelength: 668 nm                                                                                                                                                                                             |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

|                            |                                                                        |
|----------------------------|------------------------------------------------------------------------|
| <b>Neutralization</b>      | Optimal dilution of this antibody should be experimentally determined. |
| <b>Western Blot</b>        | Optimal dilution of this antibody should be experimentally determined. |
| <b>Immunocytochemistry</b> | Optimal dilution of this antibody should be experimentally determined. |

#### PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

#### BACKGROUND

Canine MCP-1 (monocyte chemotactic protein-1) is an 8 kDa member of the CC chemokine family of chemotactic factors (1, 2). It is synthesized as a 101 amino acid (aa) precursor that contains a 23 aa signal sequence and a 78 aa mature segment (3). It contains no potential N-linked glycosylation sites and is not known for any posttranslational modifications. Based on human studies, MCP-1 will primarily circulate as a monomer. Noncovalent dimers are likely to be found, however. MCP-1 activity has been localized to the N-terminus (1). Cell types known to secrete MCP-1 are considerable in number, and include keratinocytes, fibroblasts, endothelium, osteoblasts, macrophages, mast cells, smooth muscle cells and astrocytes (1, 2). In the mature MCP-1 segment, there is 82% and 83% aa identity, canine to human and porcine MCP-1, respectively. When mature canine MCP-1 is compared to (125 aa) extended rodent MCP-1, there is 55% and 56% aa identity, canine to mouse and rat MCP-1, respectively. MCP-1 has three possible receptors. The first two are CCR2 (1) and CCR11 (4). The third receptor has only been identified in mice and is called L-CCR (5). Its function is unknown. MCP-1 is best known as a chemotactic agent for mononuclear cells. It also, however, induces enzyme and cytokine release in monocytes, NK cells, and lymphocytes and histamine release by basophils (1). Additionally, it is believed to reduce IL-12 production by dendritic cells and promote a Th2 phenotype in CD4<sup>+</sup> T cells (6).

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.