

DESCRIPTION

Species Reactivity	Feline
Specificity	Detects IL-1 β /IL-1F2 in ELISAs and Western blots. In sandwich immunoassays, less than 0.1% cross-reactivity with recombinant human IL-1 β , recombinant mouse IL-1 β , and recombinant canine IL-1 β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant feline IL-1 β /IL-1F2 Ala116-Ser267 Accession # P41687
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-1 is a name that designates two pleiotropic cytokines, IL-1 α (IL-1F1) and IL-1 β (IL-1F2), which are the products of distinct genes. IL-1 α and IL-1 β are structurally related polypeptides that share approximately 27% amino acid (aa) identity in feline. Both proteins are produced by a wide variety of cells in response to inflammatory agents, infections, or microbial endotoxins. While IL-1 α and IL-1 β are regulated independently, they bind to the same receptor and exert identical biological effects. IL-1 RI binds directly to IL-1 α or IL-1 β and then associates with the IL-1 R accessory protein (IL-1 R3/IL-1 R AcP) to form a high-affinity receptor complex that is competent for signal transduction. IL-1 RII has high affinity for IL-1 β but functions as a decoy receptor and negative regulator of IL-1 β activity. IL-1ra functions as a competitive antagonist by preventing IL-1 α and IL-1 β from interacting with IL-1 RI (1-4). The feline IL-1 β cDNA encodes a 267 aa precursor. A 115 aa propeptide is cleaved intracellularly by the cysteine protease IL-1 β -converting enzyme (Caspase-1/ICE) to generate the active cytokine (5, 6). The 17 kDa mature feline IL-1 β shares 63%-78% aa sequence identity with canine, cotton rat, equine, human, mouse, porcine, rat, and rhesus IL-1 β .

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.