

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse SR-A1/MSR in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse SR-AI/MSR Trp79-Ser454 Accession # AAA39747
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The scavenger receptor (SR) family comprises a group of functionally defined membrane receptors that share the common ability to bind and internalize modified forms of Low Density Lipoproteins (mLDL) (1 - 3). Family members are classified alphabetically. The A class include four proteins: the three subtypes of SR-A (AI, AII, and AIII) that are generated by alternative splicing of the same gene, and a structurally similar protein named MARCO (4). All A class SRs are multidomain trimeric type II membrane proteins. SR-AI has an N-terminal cytoplasmic domain, a transmembrane domain, a spacer domain, an α-helical coiled coil, a collagen-like domain and a C-terminal cysteine-rich domain. SR-A is expressed by most tissue macrophages, dendritic cells and Kupffer cells. It is also highly expressed by microglia in neonatal as well as Alzheimer's Disease brains. SR-AI binds a broad range of polyanionic ligands including modified proteins (e.g. Oxidized, acetylated or maleylated LDL, Advanced glycation end-product proteins), polyribonucleotides (polyguanosine and polyinosine), polysaccharides (dextran sulfate, fucoidan), phospholipids (phosphatidylserine), bacterial products (lipopolysaccharide and lipoteichoic acid) and selected chemical compounds (silica, crocidolite asbestos). The ligand-binding region has been localized to a positively charged region in the carboxyl end of the collagen-like domain. Based on its ligand binding characteristics, SR-AI is implicated in many physiological and pathophysiological functions. Studies using SR-A knockout mouse have also suggested roles of SR-A in atherogenesis, host defense and innate immunity, acquired immune responses, macrophage adhesion, and phagocytosis of apoptotic cells (1 - 3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.