

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human sFRP-4 in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse (rm) sFRP-4 is observed, and less than 5% cross-reactivity with recombinant human (rh) sFRP-3, rh sFRP-1, and rms
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human sFRP-4 Ala22-Val346 Accession # AAC04617
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

sFRP-4, also known as DDC-4, Frp, FRPHE and Secreted Frizzled Related Protein 4, is a 50-55 kDa glycoprotein expressed in brain, kidney, lung, ovary, prostate, mammary gland, and endometrium (1, 2). This protein shows complex functions with respect to cell survival: it is up-regulated with apoptosis during ovulation (3), regulates apoptosis in chondrocytes (4), and promotes apoptosis in mammary glands when expressed in transgenic mice (5). On the other hand, sFRP-4 can also act to enhance growth as it is up-regulated in endometrial and breast carcinomas (6, 7). Since it is not detected in other carcinomas such as the ovary, colon, and pancreas, this suggest that its role in cancer is likely to be tissue dependent (6). In addition sFRP-4 is characterized as a circulating phosphaturic factor expressed by tumors associated with osteomalacia that antagonizes renal Wnt signaling (7). Of all the secreted frizzled related proteins, sFRP-4 is most closely related to sFRP-3 (1). Mouse and human sFRP-4 proteins share 92% aa identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.