

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Amnionless in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Amnionless Val20-Ala358 Accession # Q9BXJ7
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Amnionless (AMN) is an approximately 48 kDa type I transmembrane protein that is required for surface expression of the extracellular membrane-associated protein Cubilin (1-5). The two form a complex termed CUBAM (4, 5). The 453 amino acid (aa) human AMN precursor contains a 19 aa signal sequence, a 338 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 75 aa cytoplasmic domain. The ECD contains two potential N-glycosylation sites and a cysteine-rich vWFC domain. Two cytoplasmic consensus FxNPxP/F sequences for clathrin-coated pit targeting mediate endocytosis of compounds bound by Cubilin (3, 5). The ECD of human AMN shares 67%, 66%, 76%, and 78% aa identity with mouse, rat, canine and bovine AMN, respectively. AMN is present during gastrulation in the visceral endoderm and required for primitive streak formation during embryonic development in mouse (6). Transcription from an alternate start site in humans results in an isoform starting at M55 that may also function in development (7). AMN is highly expressed on polarized epithelia in the apical brush border membranes of small intestine and kidney proximal convoluted tubules (3-5, 7). Intestinal CUBAM is required for absorption of cobalamin (vitamin B12) when complexed with intrinsic factor (IF), and mutations of AMN or Cubilin cause Imersund-Grasbeck syndrome, also called megaloblastic anemia-1 (4, 5, 7). In the kidney, the CUBAM complex is thought to be important for reabsorption of proteins from filtrate, notably albumin and the molecules it carries (3, 8). Cubilin or CUBAM can be tightly associated with megalin, another endocytic receptor that contributes to stable expression of Cubilin, and to uptake of vitamin and lipoprotein complexes in the kidney and placenta (1, 9).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.