

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse HPRG in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human HPRG is observed and less than 2% cross-reactivity with recombinant mouse (rm) Cystatin A, r
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse HPRG Leu19-Lys525 Accession # BAB33094
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Mouse histidine-rich glycoprotein (HPRG) is a multidomain, monomeric, secreted, 67-75 kDa member of the cystatin superfamily of molecules (1, 2). Its name derives from the fact that 22% of its amino acids (aa) are histidine and proline. In mouse, it is synthesized as a 525 amino acid (aa) precursor that contains an 18 aa signal sequence and a 507 aa mature region (3). Five distinct domains are recognized in the mature molecule. There are two N-terminal cystatin-like modules and one His-Pro-rich region that is flanked by two Pro-rich segments (3, 4). The His-Pro-rich region contains multiple tandem repeats with a GHHPH motif, while the N- and C-termini are linked by a disulfide bond (3, 5, 6). Mouse HPRG is only 60% aa identical to human HPRG, and 79% aa identical to rat HPRG. There are multiple ligands for HPRG. These include small molecular weight molecules (metal ions; heme), hemostatic molecules (heparan sulfate; TSP; plasminogen), and immune system components (T cells; macrophages) (1, 5). About 50% of plasma plasminogen circulates bound to HPRG. Upon immobilization to cell surface tropomyosin in a Zn⁺⁺-dependent manner, it is converted to plasmin by tPA (7-9). HPRG also shows antiangiogenic activity on endothelial cells (10). Finally, it binds to cytoplasmic ligand(s) exposed during cellular necrosis, and facilitates macrophage phagocytosis (11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.