

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse sFRP-3 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human sFRP-3 (R&D Systems, Catalog # 192-SF) Ala33-Asn325 Accession # AAB51298
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Secreted Frizzled Related Protein 3 (sFRP-3) was originally identified in bovine cartilage for its chondrogenic ability. Human, mouse, chick and *Xenopus* clones have also been isolated. SFRP-3 is often referred to as FRZB, other names include Fritz, Frzb1, and FRP-3. At the amino acid sequence level, sFRP-3 is highly conserved. The human protein shares 77% identity with *Xenopus*, 92% with mouse, and 94% with bovine proteins. Human sFRP-3 is strongly expressed in the developing appendicular skeleton, and cartilage of craniofacial bones. As determined by Northern blot of adult tissues, it is strongly detected in heart and placenta, as well as the brain, skeletal muscle, kidney and pancreas.

The N-terminal portion of the human protein shows 50% amino acid identity to the corresponding region of the *Drosophila* frizzled gene product, a receptor for Wg/Wnt signals. The similarity of sFRP-3 with frizzled proteins is restricted to the N-terminal cysteine-rich domain (CRD) that contains at least ten cysteine residues with highly conserved spacing between them. SFRP-3 was subsequently shown to be a soluble antagonist of Wnt signals. It lacks all transmembrane domains of frizzled proteins but retains the ability to bind Wnts. Ectopic expression of sFRP-3 mRNA has been shown to interfere with the induction of secondary axes in *Xenopus* embryos injected with Xwnt-8 mRNA.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.