

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human and mouse IL-12 R β 2 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 5% cross-reactivity with recombinant human IL-12 R β 1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-12 R β 2 Cys28-Asn622 Accession # Q99665
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 12 (IL-12), the founding member of the IL-12 family of heterodimeric cytokines, is composed of two disulfide-linked 35 kDa and 40 kDa subunits. The 35 kDa subunit (p35) is a α -helical protein homologous to IL-6 and G-CSF. The 40 kDa subunit (p40) contains one fibronectin type III and one Ig C2-like domain, and has a high degree of structural homology to type I cytokine receptors. Whereas p35 subunit is unique to IL-12, the p40 subunit is also a subunit of IL-23. IL-12 is an essential mediator of cellular-immunity that induces T cells and natural-killer cells to produce IFN- γ . It is also required for the expansion and activation Th1 cells (1, 2).

The biological activities of IL-12 are mediated through the high-affinity receptor complex composed of the IL-12 Receptor β 1 (IL-12 R β 1) and IL-12 Receptor β 2 (IL-12 R β 2) subunits. IL-12 R β 1 is a 100 kDa protein that is also a subunit of the IL-23 receptor complex. It binds IL-12/IL-23 p40 and is associated with Tyk2. IL-12 R β 2 is a 130 kDa protein that interacts with p35 and is associated with Jak2. Both receptor subunits are type I membrane proteins that share similarities with the gp130/G-CSF R subgroup in the cytokine receptor superfamily. IL-12 R β 2 cDNA encodes a 862 amino acid (aa) residue protein with a putative 27 aa residue signal peptide that is cleaved to generate the mature protein with a 595 aa residue extracellular domain, a 24 aa residue transmembrane domain and a 216 aa residue cytoplasmic region. Human and mouse IL-12 R β 2 share 68% amino acid sequence identity. Whereas IL-12 R β 1 expression has been detected in activated T cells, NK cells and B cells, the expression of IL-12 R β 2 is more restricted. Among T cells, IL-12 R β 2 is absent on naive T cells. Activation of T cells via TCR up-regulates IL-12 R β 2 expression on human Th1 but not Th2 cells (1-4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.