

DESCRIPTION

Species Reactivity	Canine
Specificity	Detects canine IL-5 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant canine IL-5 Phe20-Ser134 Accession # Q95J76
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin-5 (IL-5) is a 40-45 kDa secreted disulfide-linked homodimeric glycoprotein that plays an important role in the differentiation, growth, and function of eosinophils. It also primes basophils for histamine and leukotriene release. In mice, IL-5 also induces the proliferation, differentiation, and immunoglobulin production of B cells especially B-1 cells that constitutively express IL-5 receptor α . IL-5 is primarily produced by CD4⁺ Th2 cells. Other cell types, including mast cells, visceral smooth muscle cells, bronchial epithelium, CD16⁺ NK cells, eosinophils and $\gamma\delta$ T cells, can also produce IL-5. Canine IL-5 is synthesized as a 134 amino acid (aa) precursor that contains a 21 aa signal sequence and a 113 aa mature segment. Mature canine IL-5 shares 62%, 66%, 85%, 84%, 58%, and 56% aa sequence identity with mature human, guinea pig, porcine, feline, mouse, and rat IL-5, respectively. The receptor for IL-5 consists of a 60 kDa ligand-binding subunit (IL-5 R α) and a 120 kDa signal-transducing subunit (β_c) (1-7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.