

## DESCRIPTION

|                           |                                                                                                                                                                                                                                                                             |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Human                                                                                                                                                                                                                                                                       |
| <b>Specificity</b>        | Detects human IL-37/IL-1F7 in direct ELISAs and Western blots. In direct ELISAs, less than 2% cross-reactivity with recombinant human (rh) FIL1δ, rhFIL1ε, rhFIL1 eta, rhIL-1α, rhIL-1β, rhIL-18, and rhIL-1 H1 is observed.                                                |
| <b>Source</b>             | Polyclonal Goat IgG                                                                                                                                                                                                                                                         |
| <b>Purification</b>       | Antigen Affinity-purified                                                                                                                                                                                                                                                   |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant human IL-37/IL-1F7<br>Lys27-Asp192<br>Accession # Q9NZH6-2                                                                                                                                                                              |
| <b>Conjugate</b>          | Alexa Fluor 700<br>Excitation Wavelength: 675-700 nm<br>Emission Wavelength: 723 nm                                                                                                                                                                                         |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

**Western Blot** Optimal dilution of this antibody should be experimentally determined.

## PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

## BACKGROUND

Human interleukin 1 family member 7 (IL-1F7), also named FIL-1Z, IL-1H4, and IL-1RP1, belongs to the IL-1 cytokine family, which currently has ten members. With the exception of IL-18 that maps to human chromosome 11, all other IL-1 family members map to the same cluster on human chromosome 2. Five alternatively spliced transcripts that arise through alternate exon usage have been described. These transcripts encode five different IL-1F7 isoforms (IL-1F7a through e also referred to as isoforms 1 through 5) that have distinct expression profiles. Polymorphism in the protein sequence of IL-1F7 isoforms also exists. Like IL-1α, IL-1β and IL-18, all of the IL-1F7 variants lack a typical signal peptide. The longest IL-1F7 transcript, referred to as IL-1F7b or IL-1F7 isoform 1, encodes a 218 amino acid (aa) residues proprotein containing a 45 aa propeptide, which is removed by caspase-1 to generate the 173 aa mature segment. Mature IL-1F7b and other IL-1F7 variants lack potential N-linked glycosylation sites. The secreted mature IL-1F7b was reported to exist as a nondisulfide linked homodimers in solution, IL-1F7 shares approximately 21%, 24%, and 30% aa sequence identity with mature IL-1α, IL-1β and IL-1ra, respectively. Mouse IL-1F7 has not been reported, but human IL-1F7 is active on mouse cells. IL-1F7b binds to IL-18 Rα with low affinity but does not exert any IL-18 agonistic or antagonistic effects. IL-1F7b also binds to the IL-18BP to enhance the antagonistic effects of IL-18BP. It has been proposed that IL-1F7b form a trimeric complex with IL-18BP and IL-18 Rβ. This complex blocks IL-18 activity by sequestering the signal transducing subunit and preventing it from participating in IL-18 signaling (1-8).

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.