

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human and mouse MIF in direct ELISAs and human, mouse and rat MIF in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse MIF Pro2-Ala115 Accession # NP_034928
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

MIF (Macrophage Inhibitory Factor; also GIF) is a 12-13 kDa member of the MIF family of proteins. Although secreted, it possesses no definitive signal sequence. MIF is widely expressed, and serves to regulate immune reactions. On macrophages and B cells, MIF binds to a CD74:CD44 complex, initiating downstream signaling. On monocytes, this promotes TNF- α production plus IFN- γ -stimulated NO production. On B cells, this promotes B cell survival. MIF also binds to leukocyte CD74:CXCR2 and CD74:CXCR4 heterodimers, initiating T cell and monocyte migration. A tautomerase activity has also been shown for MIF, but the significance of this is unclear. Mouse MIF is 114 amino acids (aa) in length. Its tautomerase activity is suggested to involve the entire length of the molecule. MIF is phosphorylated at Ser91, and an additional, singular Cys residue is covalently linked to MIF at Cys60. MIF is considered to act as a homotrimer. Full-length mouse MIF shows 90% and 99% aa sequence identity with human and rat MIF, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.