

DESCRIPTION

Species Reactivity	Canine
Specificity	Detects canine IL-12 in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant human IL-12 p35/p40 and recombinant feline IL-12 p35/p40 is observed, and less than 10% cross-reactivity with recombinant r
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant canine IL-12 p40: Ile23-Ser329; p35: Arg26-Ser222 (Asp63Glu, Tyr64Leu, Ile65Tyr, Pro66Ser) Accession # p40: Q28268; p35: Q28267
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 12 (IL-12) is the founding member of the IL-12 family of heterodimeric cytokines, which have important immunological functions. IL-12 is composed of two disulfide-linked subunits of 35 kDa and 40 kDa, respectively. The 35 kDa subunit (p35) is an α -helical protein homologous to IL-6 and G-CSF. The 40 kDa subunit (p40) contains one fibronectin type III and one Ig C2-like domain, and has a high degree of structural homology to type I cytokine receptors. Whereas p35 subunit is unique to IL-12, the p40 subunit is also utilized in IL-23. Mature canine p35 is a 197 amino acids (aa) protein that is secreted as a heterodimer linked to p40. It contains two potential N-linked glycosylation sites and shares 91%, 86%, 85%, 56%, and 55% aa sequence identity with feline, human, porcine, rat and mouse p35, respectively. Mature p40 contains 307 aa and can exist in multiple forms, including monomer, homodimer, heterodimer linked to p19 (forming IL-23), and heterodimer linked to p35 (forming IL-12). It is upregulated by substances such as LPS and CpG that activate antigen-presenting cells. Mature canine p40 shows 94%, 85%, 84%, 65%, and 65% aa sequence identity to feline, human, porcine, rat and mouse p40, respectively. Cells known to produce IL-12 include macrophages, dendritic cells, monocytes, Langerhans cells, neutrophils, and keratinocytes. The activities of IL-12 are mediated by the receptor complex composed of two type I transmembrane proteins: a 100 kDa ligand-binding subunit (IL-12 R β 1) and a 130 kDa signal transducing subunit (IL-12 R β 2). IL-12 facilitates hematopoietic stem cell proliferation, induces NK cell proliferation, promotes IgG production (in mice), and potentiates the expansion and late activation of Th1 CD4⁺ T cells (1-5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.