

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Siglec-3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human (rh) Siglec-7 and rhSiglec-9 and less than 5% cross-reactivity with recombinant mouse (rm) Siglec-F, rmSiglec-2, rhSiglec-3, rhSiglec-5, and rhSiglec-10 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse Siglec-3/CD33 Gln17-Val244 Accession # Q63994
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse Siglec-3/CD33

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Siglecs (sialic acid binding Ig-like lectins) are I-type (Ig-type) cell-surface lectins that recognize sialic acid within specific glycan structures. Mouse Siglec-3, also known as CD33 and GP67, is expressed primarily on myeloid precursors in the bone marrow and on granulocytes in peripheral blood. Two isoforms have been reported for mouse Siglec-3 that display distinct cytoplasmic domains resulting from differential mRNA splicing.