

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-12 in direct ELISAs and Western blots. In direct ELISAs, approximately 35% cross-reactivity with recombinant human (rh) FGF-13B is observed, 10% cross-reactivity with rhFGF-11 is observed and less than 1% cross-reactivity
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FGF-12 Glu67-Thr243 Accession # P61328
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast growth factor 12 (FGF-12) is a member of the FGF superfamily of molecules which contains at least 22 members. The FGFs are characterized by the presence of a core, 120 amino acid (aa) β -trefoil structure, and all apparently bind heparin (1, 2). FGF-11, -12, -13, -14, originally termed FGF homologous factors (FHF) -3, -1, -2, -4, respectively, form a subgroup within the FGF family (1, 3). Human FGF-12/FHF-1 is synthesized as a 243 aa protein. It lacks a typical signal sequence and is considered to be a cytoplasmic protein. It does, however, possess an N-terminal bipartite nuclear localization signal (NLS) at aa 11-18 and 28-38 (4-6). The 243 aa protein has at least one alternate splice form that is 181 aa in length. This is termed FGF-12B. Alternate splicing deletes the N-terminal 66 aa in FGF-12 and replaces them with four aa in FGF-12B. This substitution removes the NLS from the short form. Studies on the short form (12B) show that it cannot bind any of the common FGF receptors. This is consistent with its cytoplasmic localization. It can, however, bind to IB2 (islet brain-2), a cellular kinase scaffold protein, and voltage-gated sodium channels, suggesting FGF-12B plays an important role in intracellular signaling and ion exchange (7). Mouse and human FGF-12B differ by only one amino acid.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.