

DESCRIPTION


| | |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Species Reactivity | Human |
| Specificity | Detects human PDGF R β when phosphorylated at Y1021 in Western blots. |
| Source | Polyclonal Rabbit IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Phosphopeptide containing human PDGF R β Y1021 site |
| Formulation | Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|---------------------|----------------------------------|---------------|
| Western Blot | 0.5 μ g/mL | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | <p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

PDGF is a major serum mitogen that can exist as a homo or hetero-dimeric protein consisting of disulfide-linked PDGF-A and PDGF-B chains. The PDGF-AA, PDGF-BB and PDGF-AB isoforms have been shown to bind to two distinct cell surface PDGF receptors with different affinities. Where as PDGF R α binds all three PDGF isoforms with high affinity, PDGF R β binds PDGF-BB only with high-affinity. Both PDGF R α and PDGF R β are members of the class III subfamily of receptor tyrosine kinases (RTK) that also includes the receptors for M-CSF, SCF and Flt3 ligand. All class III RTKs are characterized by the presence of five immunoglobulin-like domains in their extracellular region and a split kinase domain in their intracellular region. PDGF binding induces receptor homo- and hetero-dimerization and signal transduction. The expression of the α and β receptors is independently regulated in various cell types. Recombinant soluble PDGF R β binds PDGF with high affinity and is potent PDGF antagonist.

References:

1. Heldin, C.H. and L. Claesson-Welsh (1994) in *Guidebook to Cytokines and Their Receptors*, Nicola, N.A. ed. Oxford University Press, New York, p. 202.