

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-1 Rrp2/IL-1 R6 in direct ELISAs and Western blots. In direct ELISAs, less than 60% with recombinant rat IL-1 Rrp2 and recombinant human IL-1 Rrp2 is observed and less than 5% cross-reactivity with recombinant mouse (rm) IL-1 R4 is ob
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-1 Rrp2/IL-1 R6 Asp22-Arg338 (Ala158Val) Accession # Q9ERS7
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-1 Rrp2, also known as IL-1 R6, is a member of the IL-1 R/TLR superfamily. Members of this superfamily have a characteristic Toll-IL-1 R (TIR) domain in their cytoplasmic region. This superfamily can be further divided into three subgroups based on extracellular region analogies. IL-1 Rrp2 belongs to Subgroup 1 which is characterized by the presence of at least one immunoglobulin-like domain (1). Mouse IL-1 Rrp2/IL-1R6 is a 70 kDa type I transmembrane glycoprotein that is synthesized as a 574 amino acid (aa) precursor. The precursor contains a 21 aa signal sequence, a 317 aa extracellular region, a 21 aa transmembrane segment, and a 215 aa cytoplasmic domain (2). The extracellular region contains three C-type Ig-like domains. Within the extracellular region, mouse IL-1 Rrp2 is 64% and 80% aa identical to human and rat IL-1 Rrp2, respectively. Expression of IL-1 Rrp2 is restricted to lung epithelium, monocytes, skin-derived keratinocytes, fibroblasts and, to a lesser extent, endothelial cells (3). Studies have demonstrated that the functional receptor for IL-1F6, IL-1F8 and IL-1F9 is a combination of IL-1 Rrp2 and IL-1 RAcP. This suggests that IL-1 Rrp2 ligands may signal in a similar fashion to IL-1 and IL-18. In each case, there is a binding receptor which, upon ligation, recruits a second receptor as a signaling component, forming an active heterodimeric receptor complex (4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.