

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-4 in direct ELISAs and Western blots. In direct ELISAs, approximately 20% cross-reactivity with recombinant human (rh) FGF-6 is observed and less than 2% cross-reactivity with bovine FGF acidic, bovine FGF basic, rhFG
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FGF-4 Ala31-Leu206 Accession # P08620
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

FGF-4, the product of a developmentally regulated gene (*hst-1*), is a member of the FGF family that is efficiently secreted. The gene for FGF-4 (also known as FGFK or K-FGF for Kaposi sarcoma-associated FGF) was initially discovered as a transforming gene by the NIH-3T3 focus formation assay using DNA derived from human tumors (including stomach and colon cancers, hepatocellular carcinomas, and Kaposi's sarcomas). FGF-4 does not seem to be expressed in normal adult tissues. However, expression of the gene is spatially and temporally regulated during embryonic development. The murine homologue of human FGF-4 has been cloned and shown to be 82% homologous to the human protein at the amino acid sequence level. Human FGF-4 has been shown to exhibit cross species activity.

In vitro, FGF-4 is mitogenic for fibroblasts and endothelial cells. FGF-4 has been shown to be a potent angiogenesis promoter *in vivo*. FGF-4 has potent transforming potential apparently through an autocrine mechanism of action. FGF-4 plays a key role in limb development and has been identified as the molecular mediator of the activities of the apical ectodermal ridge that is required for directing the outgrowth and patterning of vertebrate limbs.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.