

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Endorepellin/Perlecan in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Endorepellin/Perlecan Arg3684-Ser4391 (Ser4003Thr) Accession # P98160
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human Endorepellin/Perlecan
Immunohistochemistry	5-15 µg/mL	See Below

DATA

Immunohistochemistry



Endorepellin/Perlecan in Human Skin.

Endorepellin/Perlecan was detected in immersion fixed paraffin-embedded sections of human skin using 5 µg/mL Goat Anti-Human Endorepellin/Perlecan Antigen Affinity-purified Polyclonal Antibody (Catalog # AF2364) overnight at 4 °C. Tissue was stained with the Anti-Goat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS008) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Endorepellin is the C-terminal domain of the heparan sulfate proteoglycan perlecan, which is a modular protein found in basement membranes. Endorepellin has been found to be an inhibitor of angiogenesis that blocks endothelial cell adhesion to fibronectin and type I collagen.