

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat IL-3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 5% cross-reactivity with recombinant mouse IL-3 and less than 1% cross-reactivity with recombinant human IL-3 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant rat IL-3 Ile27-Cys169 Accession # P97688
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Rat interleukin-3 (IL-3; also multi-CSF) is a 26 kDa, variably glycosylated monomeric polypeptide that belongs to the α -helix family of hematopoietic cytokines (1, 2). IL-3 has pleiotrophic activities on a number of hematopoietic-related cells (1, 3). The rat molecule has two alternate splice forms. The first is termed IL-3 β and is synthesized as a 169 amino acid (aa) precursor that contains a 27 aa signal sequence and a 142 aa mature segment (1, 2). The second is called IL-3 α , and is identical to IL-3 β , save for a three amino acid (Tyr-Pro-Gln) deletion at positions 56-58 (1). The beta form is considered the most common form. Each form has an α -helical structure with two intrachain disulfide bonds and two potential N-linked glycosylation sites. Rat IL-3 is generally considered to be species-specific in its activity. In the mature region, rat IL-3 shares 55%, 30%, and 24% aa sequence identity with mouse, human, and bovine IL-3, respectively. Cells known to express IL-3 include connective tissue mast cells, astrocytes, microglia, megakaryocytes, eosinophils, T cells, keratinocytes and thymic epithelium.

IL-3 exerts its biological activities by binding to a 70 kDa, low affinity, ligand-binding IL-3 R α subunit, (6) which then recruits a 120 kDa, common β -chain, signal transducing subunit (7) to form a functional IL-3 receptor (1, 6, 7). Receptors for IL-3 are present on bone marrow progenitors, macrophages, mast cells, eosinophils, megakaryocytes, basophils, and various myeloid leukemic cells. IL-3 can stimulate the proliferation and differentiation of pluripotent hematopoietic stem cells as well as various lineage committed progenitors including those for neutrophils, macrophages, megakaryocytes, and erythroid cells. IL-3 can stimulate the growth of early B cells and mature macrophages, mast cells, eosinophils, basophils, and megakaryocytes. IL-3 augments the function activity of basophils, mast cells, eosinophils, and macrophages (1, 8). In combination with other molecules such as CD40L and/or IL-4, IL-3 can stimulate production of dendritic cells (1, 2, 9, 10).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.