




DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human CD155/PVR in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD155/PVR Gly27-Asn343 Accession # AAH15542
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.25 µg/mL	See Below
Immunocytochemistry	5-15 µg/mL	Immersion fixed human peripheral blood mononuclear cells
Simple Western	2.5 µg/mL	See Below
Knockout Validated	CD155/PVR is specifically detected in HEK293T human embryonic kidney parental cell line but is not detectable in CD155/PVR knockout HEK293T cell line.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CD155 [also known as PVR (poliovirus receptor) and Nectin-5 (nectin-like molecule-5)] is a 70 kDa type I transmembrane (TM) glycoprotein that is a member of the nectin-like (Nectin) family of nectin-related molecules (1). Like nectins, Nectin molecules are Ig superfamily members that contain three Ig-like extracellular domains, a TM segment, and a cytoplasmic tail. Unlike nectins, Nectin molecules cannot interact with cytoplasmic afadin (1). While Nectins serve as cell adhesion molecules, the actual functions of most Nectins are yet-to-be determined. CD155/PVR was originally isolated based on its ability to mediate polio virus attachment to host cells (2, 3). The full-length (or CD155 α isoform) is synthesized as a 417 amino acid (aa) precursor that contains a 20 aa signal sequence, a 323 aa extracellular region, a 24 aa TM segment and a 50 aa cytoplasmic tail. The extracellular region contains one N-terminal V-type and two C2-type Ig-like domains (2, 3). The V-type domain mediates polio virus binding (4). Three other isoforms exist, all of which retain the Ig-like domains. CD155 δ is transmembrane with a shortened cytoplasmic tail of 25 aa. CD155 β (352 aa) and CD155 γ (344 aa) are 60-65 kDa soluble forms that show removal of the TM segment and surrounding amino acids (2, 5). The soluble forms will bind the polio virus (due to the presence of the V-type Ig domain) but afford no protection against polio infection because of low circulating levels (5). CD155 has been demonstrated to bind vitronectin, nectin-3, and DNAM-1 (6-8). DNAM-1 binding promotes monocyte migration and NK cell killing. CD155 is expressed in all normal tissues and is highly expressed in tumor cells of epithelial and neuronal origin.

References:

1. Takai, Y. *et al.* (2003) *Cancer Sci.* **94**:655.
2. Mendelsohn, C.L. *et al.* (1989) *Cell* **56**:855.
3. Koike, H. *et al.* (1990) *EMBO J.* **9**:3217.
4. Koike, S. *et al.* (1991) *Proc. Natl. Acad. Sci. USA* **88**:4104.
5. Baur, B. *et al.* (2003) *Biochem. Biophys. Res. Commun.* **309**:175.
6. Mueller, S. and E. Wimmer (2003) *J. Biol. Chem.* **278**:31251.
7. Raymond, N. *et al.* (2004) *J. Exp. Med.* **199**:1331.
8. Lange, R. *et al.* (2001) *Virology* **285**:218.