

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CD200 R1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 5% cross-reactivity with recombinant human CD200 R is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse CD200 R1 Thr26-Pro238 Accession # BAE42266
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD200 R1, also known as OX-2 receptor, is a 90 kDa, type I transmembrane protein that belongs to the immunoglobulin superfamily. CD200 R1 is important in the regulation of myeloid cell activity (1-3). The mouse CD200 R1 cDNA encodes a 326 aa precursor that includes a 25 aa signal sequence, a 213 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 67 aa cytoplasmic domain. The ECD is composed of one Ig-like V-type domain and one Ig-like C2-type domain (4). Within the ECD, mouse CD200 R1 shares 56% and 70% aa sequence identity with human and rat CD200 R1, respectively. The ECD of mouse CD200 R1 shares 69%, 38%, 79%, and 83% aa sequence identity with the ECD of CD200 R2, 3, 4, and a CD200 R-like molecule, respectively. CD200 R1 is expressed primarily on mast cells, basophils, macrophages, and dendritic cells, (5-7) while its ligand, CD200, is widely distributed (8). Disruption of this receptor-ligand pair by knockout of the CD200 gene leads to increased macrophage number and activation, plus a predisposition to autoimmune disorders (9). Association of CD200 with CD200 R1 takes place between their respective N-terminal Ig-like domains (10). The CD200 R-like molecules may interact differently with CD200 (11, 12). The cytoplasmic domain of CD200 R1 contains two non-ITIM tyrosine residues which are required for propagating its inhibitory signals (13-15). CD200 R-like molecules, in contrast, are potentially activating receptors by means of their association with DAP12 (4, 16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.