

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human, mouse, and rat 14-3-3ζ in Western blots. In Western blots, less than 1% cross-reactivity with recombinant human 14-3-3 beta, epsilon, eta, gamma, or sigma is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human 14-3-3ζ Met1-Asn245 Accession # P63104
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.25 μg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

14-3-3 proteins are a highly conserved family of homo- and heterodimeric phosphoserine/threonine-binding proteins present in high abundance in all eukaryotic cells. 14-3-3 proteins were the first polypeptides shown to have pSer/Thr binding properties, generally recognizing the consensus sequences RSXpSXP and RXY/FXpSXP (where X is any amino acid). 14-3-3 proteins act as key regulators of intracellular signal transduction through their ability to bind specific motifs phosphorylated on serine or threonine. For example, the binding of 14-3-3 to phosphorylated BAD blocks its proapoptotic association with Bcl-XL. There are at least seven distinct 14-3-3 genes in vertebrates, alpha/beta, epsilon, eta, gamma, theta, sigma and zeta (α/β, ε, η, γ, τ/θ, σ, and ζ/δ). 14-3-3 zeta, also known as Tyrosine 3-Monooxygenase/Tryptophan 5-Monooxygenase Activation Protein, zeta isoform (YWHAZ) is a 245 amino acid, 27 kDa protein.