

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CD27 Ligand in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant mouse CD27 Ligand and recombinant human (rh) CD30 Ligand is observed, and less than 1% cross-reactivity with rhCD40 Ligand is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD27 Ligand/TNFSF7 Gln45-Pro193 Accession # AAA36175
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Human CD27 Ligand (also CD70) is a 30 kDa type 2 transmembrane glycoprotein that is a member of the TNF superfamily. It contains a 20 amino acid (aa) cytoplasmic region and a 155 aa extracellular domain that shows multiple β-strands. It is an inducible trimer that occurs on B cells, T cells, and NK cells. It regulates T cell dependant B cell differentiation into plasma cells and likely to promotes clonal expansion of T cells. Human CD27 Ligand extracellular region is 64% aa identical to mouse and rat CD27 Ligand extracellular regions.