

DESCRIPTION





| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse Serum Amyloid A1/SAA1 and mouse Serum Amyloid A2/SAA2 in direct ELISAs and Western blots. In direct ELISAs, less than 3% cross-reactivity with recombinant human (rh) SAA1, rhSAA2, recombinant mouse (rm) SAA3, and rmSAA4 is observed. |
| Source | Polyclonal Goat IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | <i>E. coli</i> -derived recombinant mouse Serum Amyloid A1 Gly20-Tyr122 Accession # P05366 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------------|---------------------------|-----------|
| Western Blot | 1 µg/mL | See Below |
| Immunohistochemistry | 5-15 µg/mL | See Below |
| Simple Western | 50 µg/mL | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

Mouse serum amyloid A protein-1 (SAA1) is a multifunctional apolipoprotein produced by hepatocytes in response to pro-inflammatory cytokines. It is secreted as a 12 kDa, 103 amino acid (aa), nonglycosylated polypeptide and circulates as part of the HDL complex. The SAA1 gene is one of three SAA genes in mouse, and, based on human, is likely to be allelic. The SAA1 gene product differs from the SAA2 gene product by only nine amino acids. In human, circulating SAA1 shows multiple proteolytically-generated isoforms, with anywhere from one to three amino acids being cleaved from either the N- or C-terminus. The same situation may exist in mouse. The amino acid sequence of mature mouse SAA1 is 72%, 72%, 67%, and 74% identical to mature human, rabbit, equine, and hamster SAA1, respectively.