

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SLITRK1 in direct ELISAs and Western blots. In these formats, approximately 5% cross-reactivity with recombinant human (rh) SLITRK2, rhSLITRK4, and rhSLITRK5 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human SLITRK1 (R&D Systems, Catalog # 3009-SK) Ala16-Ser616 Accession # Q96PX8
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SLITRK1 (Slit and Trk-like family member 1) is a charter member of the SLITRK family of proteins (1, 2). This family includes six members, all of which contain a slit-like extracellular region, and all of which (save for SLITRK1) contain a Trk-like cytoplasmic region (3). Human SLITRK1 is a type I transmembrane protein 696 amino acids (aa) in length. It contains a 17 aa signal sequence, a 605 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 53 aa cytoplasmic region. The ECD is characterized by the presence of two leucine-rich domains (LRD) that resemble those found in slit. Each LRD (aa 57 - 200; aa 374 - 517) contains six leucine-rich repeats (LRRs), followed by a Cys-rich, C-terminal LRR (aa 212 - 262; aa 529 - 579). The exact function of the LRRs is not known. Based on other LRRs, it might be assumed that they mediate protein-protein-interaction (4). There are multiple, single aa mutations in the ECD. Their significance is unclear (5). One truncating frameshift mutation is known that shows a unique 27 aa substitution after Arg 421 (6). This 448 aa form is associated with Tourette's Syndrome. As noted, the cytoplasmic region is unique among SLITRK family members. It contains no known signaling motifs, which is in contrast to SLITRK2 - 6 which show TrkA-like tyrosine phosphorylation sites. This difference is reflected in their function. SLITRK1 is a protein that is found in adult neurons of the cerebrum, thalamus and hippocampus. It induces neurite outgrowth whereas SLITRK2 - 6 block neurite extension (3). Human SLITRK1 shares 99%, 98% and 99% aa sequence identity with canine, mouse and bovine SLITRK1, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.