

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Norrin in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse Norrin is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Norrin Lys25-Ser133 Accession # Q00604
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Norrin (also called Norrie Disease Protein or NDP) is a secreted regulatory protein that remains tightly associated with the extracellular matrix (1). At least 70 mutations of Norrin have been identified in Norrie disease (X-linked retinal dysplasia) or in a minority of X-linked familial exudative vitreoretinopathy (FEVR). Both are disorders of retinal vascularization that cause blindness (2, 3). Norrin consists of disulfide-linked homodimers that oligomerize further via disulfide bridges to form higher order oligomers containing up to ten units of 12 kDa each. The cysteine-rich C-terminal domain of Norrin is homologous to von Willebrand factor, several extracellular mucin proteins, and members of the TGF-β family. Molecular modeling studies predict that Norrin assumes a cysteine-knot structure typical for the TGF-β family (4). Although Norrin is not related to Wnt family proteins, it functions like a Wnt protein in that it binds with high affinity to the receptor Frizzled-4, requires LDL receptor-related protein (LRP) as a co-receptor, and induces activation of the canonical Wnt signaling pathway (5). Norrin and Frizzled-4 are expressed at relatively low levels in tissues displaying vascular phenotypes and genetic disruption of either gene in mice gives phenotypes with marked similarities (5, 6, 7). Human Norrin shares 100%, 96%, 95%, 94% and 90% amino acid identity with rhesus macaque, canine, bovine, mouse, and chick Norrin, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.