

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse FGF-21 in direct ELISAs and Western blots. In these formats, approximately 15% cross-reactivity with recombinant human (rh) FGF-21 is observed and less than 1% cross-reactivity with rhFGF-19 and rhFGF-23 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse FGF-21 Tyr30-Ser210 Accession # Q9JJN1
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.25 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Mouse FGF-21 is a 20 kDa, secreted, nonglycosylated polypeptide that is a member of the FGF family. It is synthesized as a 210 amino acid (aa) precursor that contains a 28 aa signal sequence and a 182 aa mature region. Mature mouse FGF-21 shares 81%, 80%, 79% and 92% aa sequence identity with human, bovine, canine and rat FGF-21, respectively. FGF-21 seems to be produced by the liver and exhibits unique antidiabetogenic effects in fat.