

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human FABP5/E-FABP in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 30% cross-reactivity with recombinant mouse (rm) FABP5 is observed and less than 5% cross-reactivity with recombinant human FABP1, -2, -3, -4, -6, -7, and rmFABP9 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human FABP5/E-FABP Ala2-Glu135 Accession # Q01469
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	See Below
<b>Immunocytochemistry</b>	5-15 µg/mL	See Below
<b>Simple Western</b>	10-12.5 µg/mL	See Below


## DATA

### Western Blot




**Detection of Human FABP5/E-FABP by Western Blot.** Western blot shows lysates of human heart tissue, human brain (cerebellum) tissue, and human placenta tissue. PVDF membrane was probed with 1 µg/mL of Goat Anti-Human FABP5/E-FABP Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3077) followed by HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF017). A specific band was detected for FABP5/E-FABP at approximately 15 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

### Immunocytochemistry



**FABP5/E-FABP in HUVEC Human Umbilical Vein Endothelial Cells.** FABP5/E-FABP was detected in immersion fixed HUVEC human umbilical vein endothelial cells using Goat Anti-Human FABP5/E-FABP Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3077) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (red; Catalog # NL001) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Non-adherent Cells](#).


### Simple Western



**Detection of Human FABP5/E-FABP by Simple Western™.** Simple Western lane view shows lysates of human heart tissue and human placenta tissue, loaded at 0.2 mg/mL. A specific band was detected for FABP5/E-FABP at approximately 21 kDa (as indicated) using 10 µg/mL of Goat Anti-Human FABP5/E-FABP Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3077) followed by 1:50 dilution of HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF109). This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.



### Simple Western



**Detection of Human FABP5/E-FABP by Simple Western™.** Simple Western lane view shows lysates of HepG2 human hepatocellular carcinoma cell line and HeLa human cervical epithelial carcinoma cell line, loaded at 0.2 mg/mL. A specific band was detected for FABP5/E-FABP at approximately 21 kDa (as indicated) using 12.5 µg/mL of Goat Anti-Human FABP5/E-FABP Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3077) followed by 1:50 dilution of HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF109). This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Human FABP-5, also known as epidermal fatty acid binding protein (E-FABP), is a 15 kDa member of a cytosolic fatty acid binding protein superfamily. It is associated with keratinocytes and adipocytes and is suggested to promote fatty acid availability to enzymes, protect cell structures from fatty acid attack, and target fatty acids to nuclear transcription factors. The amino acid sequence of human FABP5 is 80%, 81% and 92% identical to that of mouse, rat and bovine FABP5, respectively.