

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Serpin A8/Angiotensinogen in direct ELISAs and Western blots. In direct ELISAs, approximately 20% cross-reactivity with recombinant mouse Serpin A8/Angiotensinogen is observed, and less than 5% cross-reactivity with recombinant human Se
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Serpin A8/Angiotensinogen Asp34-Ala485 Accession # ABM85122
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Serpin A8 (serine proteinase inhibitor-clade A #8; also angiotensinogen/ANGT) is a secreted, 52-62 kDa glycoprotein member of the clade F-subfamily, serpin superfamily of protease inhibitors. It is expressed by neurons and hepatocytes, and undergoes extracellular cleavage by renin to create a ten amino acid (aa) peptide termed Ang/angiotensin I. This inactive peptide is further cleaved by ACE on the endothelial cell membrane to create bioactive Ang II. Ang II induces vasoconstriction and aldosterone release. Mature human angiotensinogen is 452 aa in length (aa 34-485). It contains Ang I (aa 34-43) that is cleaved to create Ang II (aa 34-41). Angiotensinogen may circulate in a 200 kDa complex with major basic protein (MBP), or as part of a larger 300 kDa complex with MBP and complement C3dg. Over aa 34-485, human serpin A8 shares 61% aa identity with mouse serpin A8.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.