

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TREML2 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant mouse TREML2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human TREML2 Gly19-Ser268 Accession # Q5T2D2
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Trem-like transcript 2 protein (TLT-2; also TREML2) is a 33 kDa (unglycosylated) type I transmembrane cell surface receptor and member of the Trem family of receptor proteins. Human TLT-2 is synthesized as a 321 amino acid (aa) precursor that contains an 18 aa signal sequence, a 250 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 32 aa cytoplasmic tail (SwissProt #: Q5T2D2). Within the ECD are an Ig-like V-type domain (aa 20-121) and one potential site of N-linked glycosylation. Mature human TLT-2 shares 49% aa sequence identity with mature mouse TLT-2. Cell surface expression of TLT-2 is seen for most of the B and myeloid cell lines (1, 2). TLT-2 is also expressed constitutively on CD8⁺ T cells and induced on CD4⁺ T cells after activation (2). TLT-2 plays a role in the innate and adaptive immune responses. In 2008, Hashiguchi *et al.* discovered that the B7 family member B7-H3, which also plays important roles in immune responses, specifically bound to TLT-2. According to their research, stimulation with B7-H3 transfectants preferentially up-regulated the proliferation and IFN-γ production on CD8⁺ T cells (2). Also, transduction of TLT-2 into T cells resulted in enhanced IL-2 and IFN-γ production via interactions with B7-H3, and blockade of the B7-H3:TLT-2 pathway with mAb against B7-H3 or TLT-2 efficiently inhibited contact hypersensitivity responses (2). In 2009, however, Leitner *et al.* extensively analyzed interaction of B7-H3 with TLT-2 and found no evidence for such an interaction (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.