

#### DESCRIPTION

|                                                                                                                                                                                                         |                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b>                                                                                                                                                                               | Human/Mouse/Rat                                                                           |
| <b>Specificity</b>                                                                                                                                                                                      | Detects human, mouse, and rat PLC-γ1 in Western blots.                                    |
| <b>Source</b>                                                                                                                                                                                           | Polyclonal Goat IgG                                                                       |
| <b>Purification</b>                                                                                                                                                                                     | Antigen Affinity-purified                                                                 |
| <b>Immunogen</b>                                                                                                                                                                                        | <i>E. coli</i> -derived recombinant human PLC-γ1<br>Ser1091-Leu1290<br>Accession # P19174 |
| <b>Conjugate</b>                                                                                                                                                                                        | Alexa Fluor 405<br>Excitation Wavelength: 405 nm<br>Emission Wavelength: 421 nm           |
| <b>Formulation</b>                                                                                                                                                                                      | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide                              |
| *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |                                                                                           |

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

**Western Blot** Optimal dilution of this antibody should be experimentally determined.

#### PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

#### BACKGROUND

Phospholipase C gamma-1 (PLC-γ1) is activated by receptor tyrosine kinases in response to growth factors and hormones. Activated PLC-γ1 catalyzes the hydrolysis of phosphatidylinositol 4,5-bisphosphate to produce the second messengers inositol 1,4,5-trisphosphate (IP<sub>3</sub>) and diacylglycerol (DAG). IP<sub>3</sub> mobilizes the release of calcium while DAG activates protein kinase C. PLC-γ1 plays an important role in regulating cell proliferation and differentiation.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.