

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human DSCAM-L1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 20% cross-reactivity with recombinant human DSCAM is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human DSCAM-L1 Glu19-Lys1591 Accession # Q8TD84
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 µg/mL	Recombinant Human DSCAM-L1
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

DSCAM-L1 is a 250 kDa type I transmembrane Ig superfamily glycoprotein that is predominantly expressed in Purkinje cells and neurons of the cerebral cortex and olfactory bulb. DSCAM-L1 mediates cell adhesion by cation-independent homophilic interactions. It may function in the formation and maintenance of neural networks and play a role in neural disorders such as Gilles de la Tourette and Jacobsen syndromes. Human DSCAM-L1 contains ten Ig-like C2-type domains and six fibronectin type-III domains. An alternately spliced isoform is generated by deletion of 211 amino acids near the N-terminal. Within the extracellular domain, human DSCAM-L1 shares 98% amino acid sequence identity with mouse and rat DSCAM-L1.