

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CXCL6/GCP-2 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant mouse (rm) MIP-2, rmKC, recombinant rat (rr) CINC-2α, rmCRG-2, recombinant human (rh) NAP-2, rrCINC-2β and rhMCP-
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CXCL6/GCP-2 Val40-Asn114 Accession # P80162
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

GCP-2 (granulocyte chemotactic protein-2) also known as CXCL6, is a CXC chemokine initially isolated as a neutrophil chemoattractant from the MG-63 osteosarcoma cell line. Among human CXC chemokines, GCP-2 is most closely related to ENA-78 (78% amino acid (aa) sequence identity in the mature peptide region and 86% identity in the signal sequence). The structure and sequence of the genes for human GCP-2 and ENA-78 also exhibit close similarity suggesting the two genes may have originated from a gene duplication. LIX (LPS-induced CXC chemokine) was initially cloned as a gene induced by LPS in mouse fibroblasts. The predicted LIX protein sequence is identical to a previously purified mouse protein designated mouse GCP-2 based on its amino sequence similarity (60% sequence identity) to human GCP-2. Mouse GCP-2/LIX is also 54% identical with human ENA-78 at the amino acid sequence level.

Human GCP-2 cDNA encodes a propeptide of 114 amino acid residues with a predicted 37 aa residue signal peptide and 77 aa residue mature protein. Several forms of natural GCP-2 have been isolated from MG-63 conditioned media, indicating that GCP-2 undergoes limited processing at both the N- and C-termini. Human GCP-2 is a primary response gene whose induction by cytokines is attenuated by dexamethasone.

Human GCP-2 and mouse GCP-2/LIX have been shown to chemoattract and activate neutrophils, but not eosinophils and monocytes. It is likely that GCP-2 activities are mediated via the human or mouse CXCR2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components were employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.