

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects endogenous human, mouse and rat 11 β -HSD1 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human 11 β -HSD1 Asn24-Lys292 Accession # P28845
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Hydroxysteroid (11-beta) Dehydrogenase, type I 11-beta-hydroxysteroid dehydrogenase 1 (11 β -HSD1, HSD11B1 or HSD1) is a microsomal glycoprotein enzyme, with an apparent mass of 36 kDa, that catalyzes the conversion of the stress hormone cortisol to the inactive metabolite cortisone. 11 β -HSD1 can also catalyze the reverse reaction, the conversion of cortisone to cortisol. 11 β -HSD1 has been detected in a wide range of rat and human tissues, including lung and testis, but highest amounts are expressed in the liver. Human 11 β -HSD1 shares 79 and 77% sequence identity with mouse and rat 11 β -HSD1, respectively. There are at least 2 isoforms of 11 β -HSD. The other isoform, type II, is expressed predominantly in the kidney and placenta and catalyzes only the 11-beta-dehydrogenation reaction. High levels of cortisol can lead to visceral obesity and particular defects in the expression or function of 11 β -HSD1 have also been associated with insulin resistance and cognitive function.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.