

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human 53BP1 when phosphorylated at S25. |
| Source | Polyclonal Rabbit IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Phosphopeptide containing human 53BP1 S25 site |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|---------------------|----------------------------------|---------------|
| Western Blot | 1 µg/mL | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|---|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | <p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

The p53-binding protein 1 (53BP1) localizes to sites of damaged DNA and is phosphorylated by the ATM checkpoint kinase. 53BP1 is required for phosphorylation of other downstream ATM substrates, such as p53 and SMC-1, and is itself a substrate for the ATM kinase. ATM phosphorylates 53BP1 at S25, and this phosphorylation aids in 53BP1 nucleating a DNA damage response protein complex.