

DESCRIPTION



Species Reactivity	Human/Mouse
Specificity	Detects human and mouse GLI-1.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse GLI-1 Met1-Glu237 Accession # P47806
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Immunocytochemistry	5-15 µg/mL	See Below

DATA

<p>Western Blot</p>  <p>Detection of Human GLI-1 by Western Blot. Western blot shows lysates of HEK293 human embryonic kidney cell line transfected with human GLI-1. PVDF membrane was probed with 1 µg/mL of Goat Anti-Human/Mouse GLI-1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3455) followed by HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF017). A specific band was detected for GLI-1 at approximately 165 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p>Immunocytochemistry</p>  <p>GLI-1 in HeLa Human Cell Line. GLI-1 was detected in immersion fixed HeLa human cervical epithelial carcinoma cell line using 10 µg/mL Goat Anti-Human/Mouse GLI-1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3455) for 3 hours at room temperature. Cells were stained with the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (red; Catalog # NL001) and counterstained with DAPI (blue). View our protocol for Fluorescent ICC Staining of Cells on Coverslips.</p>
--	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

GLI-1 is a member of the Kruppel family of zinc finger proteins. GLI-1 is activated by the sonic hedgehog pathway and influences transcription of a variety of target genes by binding to the consensus site 5'CGGGTGGTC3'. GLI-1 activation leads to cellular proliferation and anti-apoptotic activities based on target genes activated.