

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Nectin-3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 50% cross-reactivity with recombinant human Nectin-3 is observed and less than 2% cross-reactivity with recombinant mouse (rm) Nectin-1 and rmNectin-4 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Nectin-3 Leu56-Asp400 Accession # Q9JLB9.1
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse Nectin-3

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Nectin-3, also known as poliovirus receptor-related 3, is a 95 kDa type I transmembrane glycoprotein belonging to the Nectin family of Ig superfamily proteins. The extracellular domains of the Nectin family form cis-homodimers that are homophilic and heterophilic trans-interacting cell adhesion molecules, which play important roles in various cell-cell junctions. Mouse Nectin-3 extracellular domain shares 96% and 94% amino acid sequence identity with that of rat and human Nectin-3, respectively.