

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human R-Spondin 3 in direct ELISAs. In direct ELISAs, less than 10% cross-reactivity with recombinant mouse R-Spondin 3 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human R-Spondin 3 Gln22-His272 Accession # Q9BXY4
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

R-Spondin 3 (RSPO3, roof plate-specific spondin 3), also called cysteine-rich and single thrombospondin domain containing-1 (Cristin 1), is an ~31 kDa secreted protein that shares ~40% amino acid (aa) identity with the other three R-Spondin family members (1, 2). All are positive modulators of Wnt/β-catenin signaling, but each has a distinct expression pattern (1-4). Like other R-spondins, R-Spondin 3 contains two adjacent cysteine-rich furin-like domains (aa 35-135) with one potential N-glycosylation site (aa 36), followed by a thrombospondin (TSP-1) motif (aa 147-207) and a region rich in basic residues (aa 211-269). Only the furin-like domains are needed for β-catenin stabilization (2). Within aa 21-209, human R-Spondin 3 shares 93%, 92%, 97%, 96% and 92% aa identity with mouse, rat, equine, bovine and canine R-Spondin 3, respectively. Potential isoforms of 279 and 297 aa diverge at aa 210 and 276, respectively (5). Mouse R-Spondin 3 is critical for development of the placental labyrinthine layer, probably by promoting VEGF expression and thus vascular development (6, 7). It is also essential for expression of the placenta-specific transcription factor, Gcm1. In the mouse embryo, R-Spondin 3 is often expressed by or located near endothelial cells (6). It is found in the roof plate, tail, somites, otic vesicles, cephalic mesoderm, truncus arteriosus, atrioventricular canal of the developing heart, and strongly but transiently in developing limbs (4, 7). R-Spondins regulate Wnt/β-catenin by competing with the Wnt antagonist DKK-1 for binding to the Wnt co-receptors LRP-6 and Kremen, reducing their DKK-1-mediated internalization (8, 9). Reports differ on whether R-Spondins bind LRP-6 directly (8-10). R-Spondin 3 has also been identified as an oncogene (11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.